

# Anaphylaxis

## Pre-module questions

1. The most important therapeutic agents used to treat anaphylaxis are:
  - a. Oxygen and diphenhydramine
  - b. Epinephrine and diphenhydramine
  - c. Albuterol and epinephrine
  - d. Oxygen and epinephrine
2. Signs and symptoms of anaphylaxis include all of the following **EXCEPT**:
  - a. Vomiting
  - b. Flushing
  - c. Bradycardia
  - d. Syncope

## Post-module questions

1. If a patient is experiencing anaphylaxis, the first step in management is:
  - a. Assessing airway, breathing and circulation
  - b. Administering epinephrine
  - c. Calling the PICU
  - d. Placing patient in recumbent position
2. Which of the following therapeutic agents is correctly paired with the symptom it can relieve?
  - a. H1-antihistamine....bronchospasm
  - b. Inhaled beta-2 agonist...airway edema
  - c. Glucocorticosteroid...acute edema



National Pediatric Night  
Float Curriculum  
Nicole Paradise Black, M.D.  
Shands Children's Hospital  
at University of Florida

- d. H1-antihistamine...urticaria

**Knowledge test questions**

1. The recommended route of administration of epinephrine in the setting of anaphylaxis is:
  - a. Intramuscular
  - b. Subcutaneous
  - c. Endotracheal
  - d. Intravenous
  
2. Which of the following scenarios does **NOT** fit the diagnostic criteria for anaphylaxis? Sudden onset:
  - a. flushing with lip and tongue swelling and dyspnea
  - b. urticaria after starting amoxicillin for acute otitis media
  - c. pruritis and hypotension after starting ceftriaxone
  - d. hypotension after exposure to latex in a patient with latex allergy



National Pediatric Night  
Float Curriculum  
Nicole Paradise Black, M.D.  
Shands Children's Hospital  
at University of Florida

### Pre-module answers

3. The most important therapeutic agents used to treat anaphylaxis are:
  - a. Oxygen and diphenhydramine
  - b. Epinephrine and diphenhydramine
  - c. Albuterol and epinephrine
  - d. Oxygen and epinephrine**
  
4. Signs and symptoms of anaphylaxis include all of the following **EXCEPT**:
  - a. Vomiting
  - b. Flushing
  - c. Bradycardia**
  - d. Syncope

### Post-module answers

1. If a patient is experiencing anaphylaxis, the first step in management is:
  - a. Assessing airway, breathing and circulation**
  - b. Administering epinephrine
  - c. Calling the PICU
  - d. Placing patient in recumbent position
  
2. Which of the following therapeutic agents is correctly paired with the symptom it can relieve?
  - a. H1-antihistamine....bronchospasm
  - b. Inhaled beta-2 agonist...airway edema
  - c. Glucocorticosteroid...acute edema
  - d. H1-antihistamine...urticaria**

### Knowledge test answers

1. The recommended route of administration of epinephrine in the setting of anaphylaxis is:
  - a. Intramuscular**
  - b. Subcutaneous



National Pediatric Night  
Float Curriculum  
Nicole Paradise Black, M.D.  
Shands Children's Hospital  
at University of Florida

- c. Endotracheal
  - d. Intravenous
2. Which of the following scenarios does **NOT** fit the diagnostic criteria for anaphylaxis? Sudden onset:
- a. flushing with lip and tongue swelling and dyspnea
  - b. urticaria after starting amoxicillin for acute otitis media**
  - c. pruritis and hypotension after starting ceftriaxone
  - d. hypotension after exposure to latex in a patient with latex allergy



National Pediatric Night  
Float Curriculum  
Nicole Paradise Black, M.D.  
Shands Children's Hospital  
at University of Florida